



VIDAS[®] NT-proBNP2

DIAGNOSTIC WORKUP OF PATIENTS WITH SUSPECTED HF

SUSPECTED HF - ACUTE ONSET OF SYMPTOMS

PATIENT ASSESSMENT BASIC WORKUP: Physical exam and history, ECG, Chest X-ray, Lab tests

VIDAS[®] NT-proBNP2

HF UNLIKELY

< 300 pg/mL

CONSIDER other DIAGNOSIS

GREY ZONE

≥ 300 pg/mL
but under age-adjusted
rule-in cut-offs

HF LIKELY

≥ 450 pg/mL	< 50 years
≥ 900 pg/mL	50-75 years
≥ 1,800 pg/mL	> 75 years

CONFIDENTLY RULE-OUT HEART FAILURE in **ACUTE SETTINGS**

NT-proBNP concentrations must be interpreted in conjunction with the patient's previous history, physical examination or any other clinical data (e.g.: medical imaging, analyses).
VIDAS[®] NT-proBNP2 Package insert, P Ponikowski; A.A. Voors European Heart Journal 2016.



VIDAS® NT-proBNP2

DIAGNOSTIC WORKUP OF PATIENTS WITH SUSPECTED HF

SUSPECTED HF - NON ACUTE ONSET OF SYMPTOMS

HF probability ASSESSMENT

1- Clinical history

2- Physical examination

3- ECG

ALL ABSENT

≥ 1 (from the 3 above)

VIDAS® NT-proBNP2

< 125 pg/mL

<75 y.o. | ≥ 125 pg/mL
≥75 y.o. | ≥ 450 pg/mL

HF UNLIKELY
CONSIDER other DIAGNOSIS

HF LIKELY

CONFIDENTLY ASSESS HEART FAILURE in **NON-ACUTE SETTINGS**

NT-proBNP concentrations must be interpreted in conjunction with the patient's previous history, physical examination or any other clinical data (e.g.: medical imaging, analyses).
VIDAS® NT-proBNP2 Package insert, P Ponikowski; A.A. Voors European Heart Journal 2016.

